

Theoretical Physics Institute
University of Minnesota

TPI-MINN-93/91-T

UMN-TH-1212/93

August 1993

Infinite renormalization of θ -term
and Jarlskog invariant for CP -violation

I.B.Khriplovich¹

Budker Institute of Nuclear Physics

630090 Novosibirsk, Russia

and

A.I.Vainshtein²

Theoretical Physics Institute

University of Minnesota, 116 Church St. SE

Minneapolis, Minnesota 55455

and

Budker Institute of Nuclear Physics

630090 Novosibirsk, Russia

Abstract

The logarithmic ultraviolet divergence in the θ -term induced by electroweak interactions in the Kobayashi-Maskawa model of CP -violation, found by Ellis and Gaillard, is discussed. We relate it to the Jarlskog CP -odd invariant of quark mass matrix. The divergence arises in 14th order in Higgs coupling as well as in 12th order plus 2nd order in $U(1)$ gauge boson coupling.

¹e-mail address: KHRIPLOVICH@INP.NSK.SU

²e-mail address: VAINSHTEIN@MNHEP.HEP.UMN.EDU

The purpose of this note is to comment on the connection between the infinite renormalization of the θ -term found by Ellis and Gaillard[1] in the Kobayashi-Maskawa (KM) model of CP -violation and the Jarlskog invariant[2] of quark mass matrix for CP -violating effects in this model.

It was first noted by Wilczek[3] that the parameter θ in the structure

$$\theta \frac{g_S^2}{32\pi^2} G_{\mu\nu}^a \tilde{G}_{\mu\nu}^a \quad (1)$$

can get infinitely renormalized due to CP -violating electroweak interactions. Then it was shown in ref.[1] that such infinite renormalization can occur only starting from 14th order in the weak coupling g_W . It looks only natural that operator (1) of dimension 4 gets infinitely (logarithmically) renormalized. Amazing is such a high order of perturbation theory where the infinite renormalization shows up.

On the other hand due to the GIM mechanism the very existence of this ultraviolet logarithmic divergence may look strange. Indeed in a renormalizable gauge the ultraviolet divergence of any loop contributing to the operator (1) cannot be stronger than logarithmic one. One may expect that the GIM mechanism leads to cancellation of logarithmic divergencies in the θ -term. Such an improvement of the ultraviolet behaviour takes place indeed. In particular the order $g_W^4 g_S^2$ correction to the θ -term turns out to be finite even in the local limit of the four-fermion interaction[4].

The crucial observation of ref.[1] is that the θ -term divergence arises in a renormalizable gauge not due to the $W^\pm$ -exchange, but due to the Higgs one. The Higgs coupling with fermions has the form

$$\phi^0 (\bar{D}_L H_D D_R + \bar{U}_R H_U^\dagger U_L) + \phi^+ (\bar{U}_L H_D D_R - \bar{U}_R H_U^\dagger D_L) + h.c. \quad (2)$$

Here ϕ^0, ϕ^+ are the Higgs field components,

$$U = \begin{pmatrix} u \\ c \\ t \end{pmatrix}, \quad D = \begin{pmatrix} d \\ s \\ b \end{pmatrix} \quad (3)$$

are the sets of up and down quarks, H_U, H_D are the Higgs coupling matrices directly related to the mass matrices M_U, M_D of U- and D-quarks:

$$H_Q = \frac{g_W}{2m_W} M_Q, \quad Q = U, D. \quad (4)$$

The coefficient $g_W/2m_W$ in eq.(4) has the well-known meaning of the inverse vacuum expectation value of the neutral Higgs field.

Let us emphasize that GIM cancellation occurs differently in the Higgs exchange and in the W -boson one (in a renormalizable gauge). In more common W case it is due to the subtraction of fermionic propagators with different masses which improves indeed ultraviolet behaviour of loops. For the Higgs exchange the fermion mass differences discussed can be organized directly as a proper combination of the Yukawa coupling constants. The difference in the mechanism can be conveniently formulated in the symmetric (unbroken) phase. In the first, W case expansion in masses can be interpreted as an emission of neutral Higgs quanta with vanishing 4-momentum. Meanwhile in the second case we deal with the exchange by virtual Higgs quanta and the integration over their 4-momenta, i.e., with true radiative corrections in the Higgs coupling.

The necessary combination of the Higgs coupling constants is fixed in fact by the CP -odd nature of the θ -term. It coincides (up to a factor) with the CP -odd invariant of the Higgs coupling matrices (mass matrices) found in ref.[2]:

$$J = \text{Im det}[H_U H_U^\dagger, H_D H_D^\dagger] = \left(\frac{g_W}{2m_W}\right)^{12} \text{Im det}[M_U M_U^\dagger, M_D M_D^\dagger]. \quad (5)$$

It is convenient to use the following representation of the mass matrices:

$$M_Q = e^{i\eta_Q} L_Q M_Q^d R_Q^\dagger \quad (6)$$

where $M_{U,D} = \text{diag}(m_{u,d}, m_{c,s}, m_{t,b})$ are positive definite diagonal mass matrices of up and down quarks, L_Q and R_Q are unitary unimodular matrices. Two more parameters in eq.(6) are the $U(1)$ phases $\eta_{U,D}$. The CKM matrix V is then

$$V = L_U L_D^\dagger \quad (7)$$

The perturbation expansion of the θ -term in the Higgs coupling (in renormalizable gauge) dictates just the form (5) of the invariant. Its explicit relation to the CP -odd invariant of the CKM matrix V is[2]

$$J = \left(\frac{g_W}{2m_W}\right)^{12} (m_t^2 - m_u^2)(m_t^2 - m_c^2)(m_c^2 - m_u^2)(m_b^2 - m_d^2)(m_b^2 - m_s^2)(m_s^2 - m_d^2) j \quad (8)$$

$$j = \text{Im } V_{cs} V_{us}^* V_{ud} V_{cd}^* \quad (9)$$

In the standard parametrization[5] $j = c_{12}c_{23}c_{13}^2 s_{12}s_{23}s_{13} \sin \delta_{13}$. Let us note that to derive eq.(8) it is convenient to use the simple relation $\text{det} C = (1/3)\text{Tr}(C^3)$ for a traceless matrix $C = [M_U M_U^\dagger, M_D M_D^\dagger]$ and the observation[2] that the imaginary part of the product $V_{ij} V_{kj}^* V_{kl} V_{il}^*$ either coincides (up to a sign) with j in eq.(9) or vanishes.

The necessity of using the combination $M_Q M_Q^\dagger$ in eq.(5) is determined by the independence on right-handed rotations R_Q . Simultaneously it guarantees the independence on $U(1)$ phases η_Q . Certainly one could use arbitrary powers of $M_Q M_Q^\dagger$ in the commutator (5) (see ref.[2]). Being interested in the lowest possible order in the Higgs coupling we are limiting ourselves just by the first power.

Let us come back now to the original problem of the possible logarithmic divergence in the induced θ -term, θ^{ind} . From the CP -odd nature of the effect it is clear that it should be proportional to the invariant (5),

$$\theta^{ind} \propto J \log M_{uv} \quad (10)$$

where M_{uv} is an ultraviolet cut-off. Eq.(10) implies immediately that the divergence might appear in 12th order in the Higgs coupling. Let us note that in the study of the logarithmic divergence (10) it is sufficient to consider the unbroken (symmetric) phase of the theory, with massless fermions and vector bosons.

Actually the divergence (10) is absent even in the 12th order. This is due to the invariance of the fermion-Higgs interaction (2) under the substitution³

$$\phi = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix} \leftrightarrow \phi^c = \begin{pmatrix} \bar{\phi}^0 \\ -\bar{\phi}^- \end{pmatrix}, \quad U_R \leftrightarrow D_R, \quad H_U \leftrightarrow H_D. \quad (11)$$

However, the r.h.s. of eq.(10) is odd under this transformation, $J \rightarrow -J$. Therefore to the 12th order the effect still vanishes.

One way to circumvent this cancellation between up and down quarks was found in ref.[1]. It is to take into account the interaction with the $U(1)$ neutral vector boson (linear combination of a photon and Z -boson) coupled to the weak hypercharge Y which is different for right-handed U and D -quarks, $Y(U_R) = 4/3, Y(D_R) = -2/3$. Then the logarithmic divergence in the θ -term arises in 14th order, 12th in the Higgs coupling and 2nd in the vector boson one:

$$\theta^{ind} \propto (g'_W)^2 [Y^2(U_R) - Y^2(D_R)] J \log M_{uv} \quad (12)$$

where g'_W is the $U(1)$ gauge coupling constant. The expression (12) is obviously invariant under the transformation (11). We would like to point out here that the effect arises also without going beyond the fermion-Higgs sector, to 14th order in the Higgs coupling:

$$\theta^{ind} \propto Tr(H_U H_U^\dagger - H_D H_D^\dagger) J \log M_{uv}. \quad (13)$$

³The invariance (11) is a leftover of a global $SU(2)_R$ symmetry exact in the limit $H_U = H_D$.

It is due to the extra Higgs exchange instead of a vector boson one.

We wish to make few comments more. Let us consider Feynman diagrams generating the effective θ -term (see fig.1). Here dashed lines denote external gluons, solid lines refer to fermions, Higgs and $U(1)$ boson exchanges are confined in rectangular blocks. Strictly speaking, the θ -term (1), being a total derivative, vanishes in the effective action after integrating over 4-coordinates. To avoid the complication one can use, e.g., the external field technique.

The induced θ -term is split into two parts due to diagrams 1a and 1b respectively

$$\theta^{ind} = \theta_a^{ind} + \theta_b^{ind}. \quad (14)$$

In fig.1a both gluons are coupled to the same fermion while in fig.1b their vertices are separated by electroweak exchanges. In fact, diagram 1a gives nothing else but the θ -term generated by the induced CP -odd phases of fermion mass matrices,

$$\theta_a^{ind} = \eta_U^{ind} + \eta_D^{ind} \quad (15)$$

where $\eta_{U,D}$ are defined by eq.(6). This is just that contribution to θ^{ind} which was discussed in ref.[1].

There are some subtle points in the calculation of diagram 1a. Namely, the last integration (over the momentum of a fermion coupled to gluons) is of infrared nature. It should be regularized by the explicit introduction of the fermion mass m_F so that this integral is dominated by momenta $p \sim m_F$.

There are no complications of the kind in θ_b^{ind} , the part of θ^{ind} due to the “irreducible” diagram 1b which is completely dominated by high momenta $p \sim M_{uv}$. So, the above analysis leading to formulae (12) and (13) is directly applicable to this contribution to θ^{ind} .

Coming back to the CP -odd part of the fermion mass operator (rectangular block in diagram 1a), it is interesting to note that this part arises already to 8th order in the Higgs coupling. For instance, CP -odd mass operator of u -quark can be shown to have the following structure:

$$\eta_u = \frac{m_u^{CP-odd}}{m_u} \propto \left(\frac{g_W}{2m_W}\right)^8 (m_t^2 - m_c^2)(m_b^2 - m_s^2)(m_s^2 - m_d^2)(m_d^2 - m_b^2) j \log M_{uv} \quad (16)$$

where j is the CP -odd invariant (9) of the CKM matrix. The derivation of eq.(16) is similar to that of eq.(8). In terms of the initial invariant J (see eq.(5)) expression (16) is singular in the mass differences:

$$\eta_u \propto \left(\frac{2m_W}{g_W}\right)^4 \frac{J \log M_{uv}}{(m_u^2 - m_t^2)(m_c^2 - m_u^2)} \quad (17)$$

It looks contradicting to our above analysis where we assumed a proportionality to J without extra singularities in the Higgs coupling. The explanation of the singularity in eq.(17) is as follows. It arises in the limit of degenerate U -quarks. But when we are dealing with degenerate states, it is only the sum over them which should be nonsingular in the parameter of degeneracy[6]. Indeed, adding up the 8th order contributions of U -quarks into θ^{ind} (see eqs.(15) and (6)) one gets zero:

$$\eta_U^{ind} = \eta_u^{ind} + \eta_c^{ind} + \eta_t^{ind} = 0. \quad (18)$$

Of course, the similar effect takes place for D -quarks. Moreover, the same cancellation persists in 10th order in the Higgs coupling where the singularity, say, of η_u^{ind} is one power less than in eq.(17). No singularity at all occurs in 12th order, $\eta_U \propto J \log M_{uv}$. However, in this order the cancellation takes place between U - and D -quarks, $\eta_U^{ind} + \eta_D^{ind} = 0$. In this way the part of θ generated by the mass matrix, θ_a^{ind} , has the same form (12) or (13) as the “irreducible” one, θ_b^{ind} .

In conclusion let us note that at any reasonable value of the ultraviolet cut-off parameter M_{uv} the discussed 14th order contribution to the θ -term is absolutely negligible numerically. The leading finite correction to the θ -term arises in the $g_W^4 g_S^2$ order[4]. It is “irreducible”, i.e. it is not generated by the CP-odd part of fermion mass operator.

Acknowledgments


We are grateful to V.A. Miransky, M.E. Pospelov and P. Ramond for stimulating discussions. The work was supported in part by Department of Energy under the grant DOE-AC02-83ER40105.

References


- [1] J. Ellis and M.K. Gaillard, Nucl.Phys. **B150** (1979) 141
- [2] C. Jarlskog, Phys.Rev.Lett. **55** (1985) 1039; Z.Phys. **C29** (1985) 491
- [3] F. Wilczek, Phys.Rev.Lett. **40** (1978) 279
- [4] I.B. Khriplovich, Phys.Lett. **B173** (1986) 193;
Yad.Fiz. **44** (1986) 1019 [Sov.J.Nucl.Phys. **44** (1986) 659]
- [5] Particle Data Group, Phys.Rev. **D45** part II (1992) III.65
- [6] T.D. Lee, M. Nauenberg, Phys.Rev. **6B** (1964) 1549

Figure captions

Fig. 1. Diagrams generating θ -term



(a)



(b)

Figure 1